









Sprouts, Salads and Ciders: The Growing Challenge of Fresh Produce-associated Foodborne Infections




Colleen Crowe, MPH
Foodborne and Diarrheal Diseases Branch
Centers for Disease Control and Prevention





Characteristics of foodborne outbreaks




Foodborne outbreaks due to fresh produce by specific etiology

	1970's	1980's	1990's	Total
<i>Salmonella</i>	4	7	18	29
Other bacterial	2	5	6	13
<i>Shigella</i> ■	-	9	2	11
Hepatitis A	1	5	4	10
<i>E. coli</i> O157:H7	-	-	9	9
Norwalk/like	1	3	1	5
<i>Giardia</i>	-	2	3	5
<i>Cyclospora</i>	-	-	5	5
Other viral	-	-	1	1
Unknown	18	34	38	90
Total	26	65	84	178

CDC

Foodborne outbreaks due to fresh produce by specific vehicle, N=175

No. of outbreaks




A growing public health problem

- Reported outbreaks per year have tripled since the 1970's
- Outbreaks have increased more than 10x in size
 - Suggesting that increase in number of outbreaks is not just due to better reporting
- More recent outbreaks are due to *Salmonella*, *E coli* O157: H7 and *Cyclospora*
- More outbreaks are due to fruit and salads



The possibilities.....



Microbe factors

- Emerging pathogens
 - - *Cyclospora*
- Recognition of acid tolerant organisms
 - *E. coli* O157

Host Factors

- **Increased consumption**
 - Per capita consumption up 23% from 1970 to 1996
 - Dietary guidelines promote increased consumption
- **New foods have become popular**
 - Sprouts, fresh juice smoothies
- **Convenience foods available**
 - Pre-cut melons, shredded lettuce




Environmental factors

- **Globalization of food supply**
- **Mass production and distribution**
- **Changes in farming and processing practices**



Complex chain for produce from farm to table



Examples of fresh-produce associated outbreaks

Sprouts ■

Unpasteurized juice



Examples of fresh produce-associated outbreaks

✓ Sprouts

-
- 20 known outbreaks worldwide since 1973
- 15 *Salmonella*, 4 *E coli*, 1 *B. cereus*
- 15/20 in N. America
- 14/15 since 1995



Salmonella serotype Stanley: All United States, Jan - May 1995

Number of isolates




Salmonella Stanley, Traceback - 1995

United States

50 cases in 6 states


9 alfalfa sprout growers

1 U.S. seed supplier




International

96% of U.S. patients
ate sprouts grown from
alfalfa seed from
shipper in Netherlands



Rapid growth of *Salmonella* Stanley during sprouting of alfalfa seeds



Jaquette et al. Appl Env Microbiol, 1996



Sprouts: prevention challenge

- **Point of contamination: Pre-sprouting**
- **Known barriers to contamination: None**
- **Potential for growth and survival: Survival on seeds and growth during sprouting**
- **Possible solutions:**
 - Improve hygiene on those few farms that grow seed for human consumption
 - Treat seeds before sprouting – irradiation? chlorine? heat?
 - High risk groups should avoid sprouts









Examples of fresh-produce associated outbreaks

✓ [■]Unpasteurized juice

- **12 known outbreaks since 1974**
- **9 from Apple juice or cider, 3 from Orange juice**
- **Organisms: *E. coli* O157, *Cryptosporidium*, *Salmonella***



Orange juice after 1999

Type	Fruit	Juice extracted	Warning label on finished product
Pasteurized/ from concentrate			No
Raw			Yes
“Fresh squeezed” unpasteurized			No



Orange juice outbreaks due to *Salmonella*

Year	Location	# ill	Product	Type of processing	Source of contamination
1995	Theme park A Florida	62	“Fresh squeezed” juice	Raw/ untreated	Pre-harvest or processing
1999	20 states Canada	423	“Fresh squeezed” smoothies	Pre-extraction kill step	Production or transport
2000	8 states	88	“Fresh squeezed juice”, free samples	Pre-extraction kill step	Unknown



Unpasteurized juice: prevention challenge

- Point of contamination: Preharvest (irrigation water, animal feces), during transport or processing
- Known barriers to contamination: Heat
- Potential for growth and survival: Survival below pH of 4.0 and with refrigeration
- Possible solutions:
 - Pasteurization
 - Alternative terminal kill step
 - Clear warning labels for all unpasteurized juice




Imported or Domestic

Does it matter?




E. coli O157:H7 Infections, Illinois and Connecticut, May - June, 1996



E. coli O157:H7, Illinois and Connecticut, 1996 Mesclun mix

- Traceback implicated one grower in California
- Results of follow-up investigation
 - Lettuce grown near cattle ranches
 - Wash water recycled
 - Processing shed open to environment

Multistate outbreak of *S. Newport* infections, United States, 1999



N = 78



Traceback investigation

Mangos purchased in grocery stores in:

- Pennsylvania
- Massachusetts
- Colorado





The prevention challenge

- Washing by consumers only reduces contamination by 10^1 or 10^2 organisms
 - Surface is complex and difficult to clean
 - Pathogens may adhere tightly
 - Contamination may be internal
- Once surface integrity is broken, growth can be rapid
 - Refrigeration often not standard practice
 - "Pre processing" means more growth opportunities
 - Increasing time between "pre processing" and consumption

Conclusions

- Fresh produce-associated outbreaks have increased in past 3 decades
- There are many food and pathogen combinations
- There are many possible points of contamination
 - Harvesting, initial processing, distribution and final processing
- Domestic and imported produce have caused outbreaks
- Contamination is easier to prevent than to wash off
- Additional barriers or prevention steps are needed before, during, and after processing



New prevention measures for fresh produce as a result of outbreak investigations

- **Sprouts**
 - Interim recommendations for seed treatment
 - Research being actively conducted on possible pre-sprouting control steps (e.g. chlorination of seeds)
- **Juice**
 - New labeling regulations
 - Some raw juice makers switched to pasteurization
- **Lettuce**
 - Lettuce industry undertaking steps to improve sanitation during processing
- **Mangos**
 - Farm in Brazil now has in-line chlorination
 - Better sanitation being promoted on mango farms worldwide



Challenges for the future

- **Public Health officials**
 - Continue to identify problems
 - Early detection, reporting, and investigation
- **Regulatory agencies and industry**
 - Identify safe production standards and implement them



Enhanced Surveillance through MDP Program

- **Planned Sample Design**
 - Likely representativeness
 - Provide information to explain sporadic cases
- **Serotyping of isolates**
- **Enhanced research on pathogen resistance**

CDC's Use of MDP Program Results

- May provide possible links between foodborne illness and produce sources
 - Use information in conduct of outbreak investigations
 - Use information in conduct of sporadic disease case-control studies
- Provide for the advancement in laboratory methods in food testing

Acknowledgements

Cindy Friedman
Sumathi Sivapalasingham
Josefa Rangel
Ellen Steinberg
Barbara Malton
Larry Slutsker
John Brooks
Jono Mermin
Betsy Hilborn
Kim Cook
Paul Mead
Patricia Griffin
Robert Tauxe
PulseNet Team
FDA
Many State Health departments

